

DentalPractice

* Abdelsalam Issa

True Or False:

- 1- The submandibular gland occupies most of the digastric or submandibular triangle ().
- 2- The sternocleidomastoid muscle serves as the boundary between the anterior and posterior triangles of the neck. ()
- 3- The infraorbital nerve is a branch of the maxillary nerve ().
- 4- The facial nerve leaves the skull through the stylomastoid foramen ().
- 5- Procaine is a vasodilator ().
- 6- Penicillin is effective against gram positive organisms ().
- 7- The body cells most susceptible to fluoride are the odontoblasts ().
- 8- The acute lethal dosage of fluoride is about 35 mg/ kg. of body weight ().
- 9- The characteristic clinical feature of a chronic periapical abscess is the absence of clinical symptoms ().
- 10- Lymphocytes are found in dental granulomas ().
- 11- Noma is the synonym for gangrenous stomatitis ().
- 12- The presence of ectopic sebaceous glands in oral mucous membrane is referred to Fordy's disease ().
- 13- The organism most commonly found in osteomyelitis staphylococcus aureus ().
- 14- The most common malignancy found in the oral cavity is the squamous cell carcinoma ().
- 15- Gingival hemorrhage may accompany leukemia because of an associated thrombocytopenia ().
- 16- The chemical disintegration of enamel is referred to erosion ().
- 17- Dens in dente occur most commonly in the maxillary lateral incisors ().
- 18- A pear - shaped radiolucency causing displacement of the roots of vital maxillary lateral and canine is characteristic of the globulomaxillary cyst ().
- 19- Features of familial multiple neurofibromas with café-au- lait spots of the skin are typical of von recklinghausen's disease of skin ().
- 20- koplik's spot are an early intraoral manifestation of rubella ().
- 21- Features of multiple bone radiolucencies, hypercalcemia, hypophosphatemia, and loss of lamin dura are indicative of hyperparathyroidism ().
- 22- The melkersson- Rosenthal syndrome is characterized by facial paralysis, cheilitis granulomatosa, and scrotal tongue ().
- 23- White, interlacting lines (striae of wickham) on the leuccal mucosa are a characteristic clinical feature of lichen planus ().
- 24- Tumors metastatic to the jaw bones are most common to be found in the posterior molar region ().
- 25- The best laboratory test to use in the diagnosis of lupus vulgaris in the oral cavity is biopsy ().
- 26- The most common benign tumor occurring in the oral cavity is fibroma ().
- 27- In the ideal developlasty in preparation of t a mouth for dentures the hight of the ridge should be equal to the width ().
- 28- Peutz- jeghers syndrome is characterized by intestinal polyposis, familial history of the disease and brown pigmentation of the oral mucosa ().
- 29- The most common bacterium responsible for subacute bacterial endocarditis after dental extraction is streptococcus viridans ().
- 30- Epinephrine in concentration of 1: 100,000 is considered edequate as a vasoconstrictor in lidocaine solutions for all procedures ().

* D.D.S
M.F. Surgeon